

A SHORT PROOF OF THE KHUKHRO–MAKARENKO THEOREM ON LARGE CHARACTERISTIC SUBGROUPS WITH LAWS

Anton A. Klyachko and Yulia B. Melnikova

Faculty of Mechanics and Mathematics, Moscow State University

Moscow 119992, Leninskie gory, MSU

klyachko@mech.math.msu.su

yuliamel@mail.ru

Dedicated to A. L. Shmelkin on the occasion of his seventieth birthday

We give a short proof and some strengthening of the Khukhro–Makarenko theorem that each group virtually satisfying an outer commutator identity contains a finite-index characteristic subgroup satisfying this identity. An estimate for the index of this characteristic subgroup is obtained.

Key words: characteristic subgroups, outer commutator identities.

Suppose that G is a group and H is its subgroup of finite index. Textbooks in group theory (see, e.g., [KaM82]) contain some simple facts allowing us to find a finite-index subgroup in G which is similar to, but better than H . In particular,

- H contains a normal finite-index subgroup of G (whose index divides $|G:H|!$);
- if G is finitely generated, then H contains a finite-index subgroup which is fully characteristic (and even verbal) in G ;
- if H is abelian, then G has a characteristic abelian subgroup of finite index.

Recently, the last statement was substantially generalised.

Khukhro–Makarenko Theorem ([KhM07], see also [MKh07]). *If a group G has a finite-index subgroup satisfying an outer (multilinear, in authors' original terminology) commutator identity, then G also has a finite-index characteristic subgroup satisfying this identity.*

Examples of outer commutator identities are n -step nilpotency or solvability. The general definition is as follows. Let $F(x_1, x_2, \dots)$ be a free group of countable rank. An outer commutator of weight 1 is a generator x_i . An outer commutator of weight $t > 1$ is a word of the form $w(x_1, \dots, x_t) = [u(x_1, \dots, x_r), v(x_{r+1}, \dots, x_t)]$, where u and v are outer commutators of weights r and $t - r$, respectively. Informally, an outer commutator of weight t is an expression $[x_1, x_2, \dots, x_t]$ with some arrangement of brackets. An outer commutator identity of weight t is an identity $w(x_1, \dots, x_t) = 1$ whose left-hand side is an outer commutator of weight t .

The following proof of the Khukhro–Makarenko theorem is significantly simpler and shorter than the original argument.

Suppose that $H_1, \dots, H_t$ are normal subgroups of a group G and $w(x_1, \dots, x_t)$ is an outer commutator. Then

- 1) the subgroup $w(H_1, \dots, H_t) \stackrel{\text{def}}{=} \langle w(h_1, \dots, h_t) ; h_i \in H_i \rangle$ is normal in G ;
- 2) $w(G, \dots, G) = 1$ if and only if G satisfies the identity $w(x_1, \dots, x_t) = 1$;
- 3) $w(H_1, \dots, H_t) = [u(H_1, \dots, H_r), v(H_{r+1}, \dots, H_t)]$ if $w(x_1, \dots, x_t) = [u(x_1, \dots, x_r), v(x_{r+1}, \dots, x_t)]$;
- 4) $w(H_1, \dots, H_{i-1}, \prod_{N \in \mathcal{N}} N, H_{i+1}, \dots, H_t) = \prod_{N \in \mathcal{N}} w(H_1, \dots, H_{i-1}, N, H_{i+1}, \dots, H_t)$

for any family $\mathcal{N}$ of normal subgroups of G .

These property are almost obvious and can be easily verified by induction.

Lemma. *If $w(x_1, \dots, x_t)$ is an outer commutator, m is a positive integer, G is a group, and $\mathcal{N}$ is a family of its normal subgroups such that*

$$w(\underbrace{N, N, \dots, N}_{m \text{ times}}, G, G, \dots, G) = 1 \quad \text{for all } N \in \mathcal{N},$$

then

$$w(\underbrace{\hat{N}, \hat{N}, \dots, \hat{N}}_{m-1 \text{ times}}, \hat{G}, \hat{G}, \dots, \hat{G}) = 1, \quad \text{where } \hat{N} = \bigcap_{N \in \mathcal{N}} N \text{ and } \hat{G} = \prod_{N \in \mathcal{N}} N.$$

Proof.

$$w(\underbrace{\hat{N}, \hat{N}, \dots, \hat{N}}_{m-1 \text{ times}}, \hat{G}, \hat{G}, \dots, \hat{G}) = w(\underbrace{\hat{N}, \hat{N}, \dots, \hat{N}}_{m-1 \text{ times}}, \prod_{N \in \mathcal{N}} N, \hat{G}, \dots, \hat{G}) = \prod_{N \in \mathcal{N}} w(\underbrace{\hat{N}, \hat{N}, \dots, \hat{N}}_{m-1 \text{ times}}, N, \hat{G}, \dots, \hat{G}).$$

But $\widehat{N} \subseteq N$ and $\widehat{G} \subseteq G$, therefore, each factor of the last product is contained in the group

$$w(\underbrace{N, N, \dots, N}_{m \text{ times}}, G, G, \dots, G), \quad \text{which is trivial by assumption.}$$

As a corollary, we obtain a strengthened version of the Khukhro–Makarenko theorem with an explicit bound for the index.

Theorem. *If a group G contains a finite-index subgroup N satisfying an outer commutator identity $w(x_1, \dots, x_t) = 1$, then G contains a finite-index subgroup H satisfying the same identity which is characteristic and even invariant under all surjective endomorphisms. In addition,*

$$\log_2 |G : H| \leq f^{t-1}(\log_2 |G : N|) \quad \text{if the subgroup } N \text{ is normal,} \quad (1)$$

and, therefore, $\log_2 |G : H| \leq f^{t-1}(\log_2 |G : N|)$ in the general case, where $f^k(x)$ is the k -th iteration of the function $f(x) = x(x+1)$.

Proof. For simplicity, we give a construction of a characteristic subgroup. To obtain a subgroup invariant under all surjective endomorphisms, one should replace all automorphisms by surjective endomorphisms in the argument below.

Consider the subgroup $G_1 = \prod_{\varphi \in \text{Aut } G} \varphi(N)$. This subgroup is characteristic, and $|G : G_1| \leq |G : N|$. Clearly, G_1 is a product of at most $\log_2 |G : N| + 1$ automorphic images of N (because the chain $N \subseteq N\varphi_1(N) \subseteq N\varphi_1(N)\varphi_2(N) \subseteq \dots$ cannot have more than $\log_2 |G : N| + 1$ different subgroups). Thus,

$$G_1 = \prod_{k=0}^{p_1} \varphi'_k(N), \quad \text{where } \varphi'_k \in \text{Aut } G \text{ and } p_1 \leq l_0 \stackrel{\text{def}}{=} \log_2 |G : N|.$$

Now, consider the subgroup $N_1 = \bigcap_{k=0}^{p_1} \varphi'_k(N)$. The index of an intersection of subgroups does not exceed the product of their indices (see, e.g., [KaM82]); hence,

$$l_1 \stackrel{\text{def}}{=} \log_2 |G : N_1| \leq \log_2 (|G : N|^{p_1+1}) = (p_1 + 1)l_0 \leq (l_0 + 1)l_0 = f(l_0).$$

By Lemma, we have

$$w(N_1, \dots, N_1, G_1) = 1.$$

The next step is to consider the subgroups

$$G_2 = \prod_{\varphi \in \text{Aut } G_1} \varphi(N_1) = \prod_{k=0}^{p_2} \varphi''_k(N_1) \quad \text{and} \quad N_2 = \bigcap_{k=0}^{p_2} \varphi''_k(N_1), \quad \text{where } \varphi''_k \in \text{Aut } G_1 \text{ and } p_2 \leq \log_2 |G : N_1| = l_1 \leq f(l_0).$$

Clearly, G_2 is characteristic in G (and even in G_1),

$$\log_2 |G : G_2| \leq \log_2 |G : N_1| = l_1 \leq f(l_0), \quad \text{and} \quad l_2 \stackrel{\text{def}}{=} \log_2 |G : N_2| \leq \log_2 (|G : N_1|^{p_2+1}) = (p_2 + 1)l_1 \leq f(l_1) \leq f(f(l_0)).$$

By Lemma, we have

$$w(N_2, \dots, N_2, G_2, G_2) = 1.$$

Continuing in the same manner, at the t -th step, we obtain a characteristic subgroup

$$G_t = \prod_{\varphi \in \text{Aut } G_{t-1}} \varphi(N_{t-1}) = \prod_{k=0}^{p_t} \varphi_k^{(t)}(N_{t-1}), \quad \text{where } \varphi_k^{(t)} \in \text{Aut } G_{t-1},$$

such that

$$w(G_t, \dots, G_t) = 1 \quad \text{and} \quad \log_2 |G : G_t| \leq \log_2 |G : N_{t-1}| = l_{t-1} \leq f(l_{t-2}) \leq f(f(l_{t-3})) \leq \dots \leq f^{t-1}(l_0).$$

The subgroup $H = G_t$ is as required, which proves the theorem.

Remark. In papers [KhM07] and [MKh07], there is no explicit bound, but it is mentioned that an estimate can be obtained from the proof. According to our calculations, inequality (1) is better.

REFERENCES

- [KaM82] Kargapolov M.I., Merzlyakov Yu.I. Fundamentals of group theory. Moscow: “Nauka”, 1982.
- [MKh07] Makarenko N.Yu., Khukhro E.I. Large characteristic subgroups satisfying multilinear commutator identities // Dokl. Akad. Nauk. 2007. V.412. no. 5. P.594–596.
- [KhM07] Khukhro E.I., Makarenko N.Yu. Large characteristic subgroups satisfying multilinear commutator identities // J. London Math. Soc. 2007. V.75. no.3, P.635–646.